



Brampton Diagnostic Services Immunochemistry Referral Testing Services

Direct Antiglobulin Test (DAT)

ON_REF-03

TEST DESCRIPTION

This test is performed to determine whether a patient's red blood cells have been sensitized in vivo with immunoglobulin, complement, or both. The DAT is used most commonly to investigate possible hemolytic transfusion reactions, Hemolytic Disease of the Fetus and Newborn (HDFN), autoimmune hemolytic anemia, and drug-induced immune hemolysis. The clinical significance of a DAT result should take into consideration the patient's clinical history, diagnoses, and other laboratory test results.

SPECIMEN AND REQUISITION REQUIREMENTS

Specimen(s)

- **Adults and Children:** One full (1) (preferably two (2)) 5-7 ml EDTA (lavender) tubes mixed thoroughly by gentle agitation. Within 14 days of collection.
- **Infant:** 250 microlitres minimum EDTA or microtainer tube mixed thoroughly by gentle agitation
- Label specimen with the required minimum information: patient's last name, first name, middle name, PHN or unique identifier and date of collection.

Complete Requisition (must include)

- Patient's last name, first name, middle name, PHN or unique identifier and date of birth
- Facility name, complete address, phone and fax number
- Phlebotomist ID information
- Date of collection

Requisition(s)

- *Request for Serological Investigation (F800437)*
- *Demande D'Investigation Serologique (F800850)*

PRE-SHIPING STORAGE

Recommended Refrigeration 1-10°C.

SHIPPING INSTRUCTIONS

Submit samples as soon as possible after collection.

Shipping

- Ship in a container that will maintain temperature 4°C to 25°C
- Select shipping method for container to arrive at testing site within 48 hours.
- Sample must be received at testing site within 14 days of collection.

Note: *Protect from freezing.*

SEND TO

Canadian Blood Services
Brampton Diagnostic Services
100 Parkshore Drive
Brampton, ON, L6T 5M1
Tel: 905-494-5295
Fax: 905-494-8131